



Ministero della Salute

Dipartimento per la sanità pubblica veterinaria,
la nutrizione e la sicurezza degli alimenti

Direzione Generale

della Sanità animale e del Farmaco veterinario
Ufficio II - Sanità animale ed anagrafi: profilassi
pianificate e zoonosi, anagrafi degli animali

N..... DGSA.II

Risposta al Foglio del

N.

OGGETTO: **Malattia emorragica virale del coniglio – segnalazione di nuova variante genetica ad inalterata alta patogenicità**

Roma



85496306

Regioni e province autonome

Assessorato alla Sanità

Servizio veterinario

Loro Sedi

Associazioni allevatori cunicoli

Loro sedi

e, p.c.

Centro nazionale di referenza per le Malattie virali dei lagomorfi

Brescia

Istituti zooprofilattici sperimentali (I.Z.S.)

Loro sedi

Ufficio VIII PIF e UVAC della D.G.S.A.

Sede

Con riferimento all'oggetto si informano gli Enti e le Associazioni in indirizzo che si è appreso da un articolo su un organo di stampa scientifica internazionale (che ad ogni buon fine si allega in copia) della comparsa nel nord della Francia dalla fine dell'estate 2010, di una nuova variante genetica del virus della malattia emorragica virale del coniglio (RHD o MEV), che mantiene l'usuale elevata patogenicità.

Dai pochi dati preliminari presentati nell'articolo, la cui interpretazione è confermata dal Centro di Referenza di Brescia che legge per conoscenza, si evince che i vaccini attualmente disponibili per la prevenzione della malattia potrebbero avere un effetto protettivo significativamente ridotto, in considerazione dell'ampia differenza genetica e forse antigenica dei ceppi utilizzati.

In attesa di nuovi e utili risultati dagli studi in corso dei ricercatori francesi e possibilmente anche dal Centro di Referenza che ha richiesto l'acquisizione della nuova variante quale Laboratorio di Referenza OIE per RHD, si sollecitano i Servizi veterinari localmente competenti a intensificare le misure di sorveglianza sugli allevamenti cunicoli, consigliando l'adozione di appropriate misure igienico-sanitarie e di biosicurezza. Particolare attenzione dovrà essere posta a quegli allevamenti che praticano con regolarità la profilassi vaccinale, soprattutto nel caso di mancata o parziale protezione, considerato che per tale malattia la profilassi vaccinale è ritenuta tra i presidi più concreti per prevenirne la comparsa. Per tali allevamenti si chiede di segnalare con prontezza all'I.Z.S competente per territorio, tutti gli eventuali casi di malattia conclamata con evidente mortalità in particolare fra gli animali riproduttori (o comunque vaccinati). L'I.Z.S. di cui sopra dopo aver effettuato le indagini diagnostiche che gli consentano di sospettare la malattia deve contattare immediatamente il Centro di referenza nazionale di Brescia per concordare i tempi e le modalità di invio dei campioni per la conferma. Giova ricordare che l'RHD rientra tra quelle di cui all'art. 1 del Regolamento di Polizia Veterinaria e che pertanto per essa si adottano le norme previste in quest'ultimo.

In attesa di nuove informazioni e conferme si invitano codesti Enti ed Associazioni a dare massima diffusione della presente affinché, ognuno per gli aspetti di propria competenza, possa attuare le misure più idonee e segnalare immediatamente qualsiasi sospetto di presenza della nuova variante di RHD.

Si ringrazia per la cortese attenzione.

IL DIRETTORE GENERALE

Letta F.



FIG 1:
Macroscopic lesions observed in a dead domestic rabbit from a rabbitry. The lesions are specific to rabbit haemorrhagic disease: haemorrhages in lungs, heart and kidneys, pale and friable liver, and congested and enlarged spleen were all seen

Some cases of RHD had been reported in rabbitries in north-western France, but since October 2010 this number has increased. Mortalities occurred in RHDV-vaccinated and non-vaccinated does and fattening rabbits. Emergency vaccination stopped the mortalities but in seven to 15 days, compared with seven to nine days when a 'classical' RHDV is responsible for the disease. Up to the end of January 2011, approximately 60 rabbitries had been affected by this virus in north-western and northern France. During the same period, outbreaks were reported by the SAGIR network (Lamarque and others 2000) in numerous wild rabbit populations in the same areas. Estimated mortalities are unusual, reaching 80 to 90 per cent. They are similar to those observed when RHD emerged and spread in France in native populations at the end of the 1980s.

We analysed three cases of RHD in rabbitries in north-western France, which occurred in mid-October. Postmortem examinations of dead rabbits (Fig 1) and microscopic examinations of organs (lung, liver and kidney) revealed typical RHD lesions (Plassart and others 1992).

Samples of liver from domestic and wild rabbits were frozen for virological analyses. For the screening of RHDV-positive samples, we amplified and sequenced a region located in the N-terminal of the gene encoding the capsid protein VP60 as previously described by Le Gall-Reculé and others (2011). All the samples were positive. The DNA sequences were closely related (98.5 to 100 per cent homology) and the protein sequences were similar, implying that the same virus was responsible for the clinical cases. This virus is related to, but highly distinct from, RHDV and antigenic variant RHDVa strains currently circulating in France (the average homology is only 85.7 per cent). The virus constitutes a new variant of RHDV.

To estimate its phylogenetic relationship with known rabbit lagoviruses, we performed phylogenetic analyses with all the nucleic sequences available in databases, including the non-pathogenic strains characterised in Italy (RCV) and recently in

Virology

Detection of a new variant of rabbit haemorrhagic disease virus in France

WE wish to report the detection of a new variant of rabbit haemorrhagic disease virus (RHDV) (Lagovirus, Caliciviridae) which is circulating in France and has been causing high mortality in domestic and wild rabbit populations since the end of the summer of 2010.

Rabbit haemorrhagic disease (RHD) is a highly infectious and fatal disease of the European rabbit (*Oryctolagus cuniculus*). Although the development of efficient commercial vaccines has made it possible to control the disease in the rabbit industry, RHD still threatens non-vaccinated wild populations.

Letters




FIG 2: Phylogenetic tree using the minimum evolution method derived for partial VP60 nucleotide (nt) sequences (nt 1188 to 1541) of 93 rabbit haemorrhagic disease virus (RHDV) and 32 antigenic variant RHDVa pathogenic strains, the Ashington strain, the low pathogenic MRCV strain and the non-pathogenic 06-11, RCV and 36 RCV-A1 strains. Bootstrap values greater than 75 per cent (for 1000 replicates) are given in italics before each node. The French reference European brown hare syndrome virus strain EBHSV-GD was used as an outgroup to root the tree. The RHDV, RHDVa and RCV-A1 branches were collapsed to highlight the main genogroups

Australia (RCV-A1) and France (06-11), as well as the assumed pathogenic Ashington strain from the UK and the low pathogenic strain from the USA (MRCV). For this purpose, we amplified and sequenced a region located in the C-terminal of the VP60 gene. Irrespective of the phylogenetic method used, analysis revealed that the RHDV variant formed a new genetic group, relatively distant from pathogenic and non-pathogenic lagoviruses described until now

(Fig 2). However, this group is more closely related to RHDV and RHDV-related viruses than to the Australian non-pathogenic strain that constitutes a new member of the genus *Lagovirus* (Strive and others 2009).

An initial experimental study is underway to confirm macroscopic and microscopic lesions of the new variant and to check the protection conferred by current commercial vaccines. Indeed, the mortalities reported in vaccinated rabbits suggest that vaccines may give only partial protection. A modified vaccination schedule or a new RHDV vaccine may be necessary to protect rabbits against this new variant. Other experimental studies will be undertaken specifically to determine the pathogenesis of the virus in very young rabbits. In addition, acquisition of complete molecular data should make it possible to determine the phylogenetic relationships with other lagoviruses and to understand the evolution of RHDV.

G. Le Gall-Reculé, F. Zwingelstein,
French Agency for Food, Environmental and Occupational Health & Safety (Anses), Avian and Rabbit Virology, Immunology and Parasitology Unit, Zoopôle-Les Croix, BP 53, 22440 Ploufragan, France
e-mail: ghislaine.legall-recule@anses.fr
S. Boucher, Labovet Conseil, rue Olivier de Serres, BP 539, 85505 Les Herbiers cedex, France
B. Le Normand, Clinique Vétérinaire des Marches de Bretagne, 47 boulevard Leclerc, 35460 St-Brice-en-Cogles, France
G. Plassiart, Laboratoire d'Anatomie Pathologique Vétérinaire de Metz, 11 avenue Leclerc de Hautecloque, BP 20317, 57006 Metz cedex 01, France
Y. Portejoie, Anjou Laboratoire, 18 boulevard Lavoisier, BP 20943, 49009 Angers cedex 01, France

A. Decors, National Hunting and Wildlife Agency (ONCFS), Department of Studies and Research, 5 rue de Saint-Thibaut, Saint Benoist, BP 20, 78612 Le Perray-en-Yvelines cedex, France

S. Bertagnoli, J.-L. Guérin, INRA, UMR1125, and Université de Toulouse, ENVT, 23 chemin des Capelles, 31076 Toulouse, France

S. Marchandeau, National Hunting and Wildlife Agency (ONCFS), Department of Studies and Research, 39 boulevard Albert Einstein, CS 42355, 44323 Nantes cedex 3, France

References

- LE GALL-RECULE, G., ZWINGELSTEIN, F., FAGES, M-P., BERTAGNOLI, S., GELFI, J., AUBINEAU, J., ROOBROUCK, A., BOTTI, G., LAVAZZA, A. & MARCHANDEAU, S. (2011) Characterisation of a non-pathogenic and non-protective infectious rabbit lagovirus related to RHDV. *Virology* 410, 395-402
- LAMARQUE, F., HATIER, C., ARTOIS, M., BERNY, P. & DIEDLER, C. (2000) Le réseau SACIR, réseau national de suivi sanitaire de la faune sauvage française. *Epidémiologie et Santé Animale* 37, 21-30
- PLASSIART, G., CUELH, J-F., CANIERE J-P., WANG, B., ANDRE-FONTAINE, G. & WYERS, M. (1992) Hematological parameters and visceral lesions relationships in rabbit viral hemorrhagic disease. *Journal of Veterinary Medicine Series B* 39, 443-453
- STRIVE, T., WRIGHT, J. D. & ROBINSON, A. J. (2009) Identification and partial characterization of a new lagovirus in Australian wild rabbits. *Virology* 384, 97-105

doi: 10.1136/vr.d697



Detection of a new variant of rabbit haemorrhagic disease virus in France

G. Le Gall-Reculé, F. Zwingelstein, S. Boucher, et al.

Veterinary Record 2011 168: 137-138
doi: 10.1136/vr.d697

Updated information and services can be found at:
<http://veterinaryrecord.bmjjournals.com/content/168/5/137.2.full.html>

These include:

References

This article cites 4 articles

<http://veterinaryrecord.bmjjournals.com/content/168/5/137.2.full.html#ref-list-1>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:
<http://group.bmjjournals.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmjjournals.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmjjournals.com/subscribe/>